



Hybrid Closed-Loop Insulin Delivery in an Adolescent with Recurrent Diabetic Ketoacidosis: A Narrative Review of Technology, Adherence and Family Support

Katarzyna Zielinska^{1*}, Lukas Bauer² and Emma Nieminen³

¹Department of Paediatric Endocrinology, Pomeranian University Hospital, Gdansk, Poland

²Department of Diabetes and Endocrinology, Styrian Institute of Clinical Medicine, Graz, Austria

³Department of Paediatrics, Ostrobothnian Medical Centre, Vaasa, Finland

***Correspondence to:** Katarzyna Zielinska, Department of Paediatric Endocrinology, Pomeranian University Hospital, Gdansk, Poland

Received: 22 June 2024; **Accepted:** 31 October 2024; **Published:** 27 December 2024

Copyright: © 2024 KZ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Adolescence is where type 1 diabetes control reliably deteriorates, and recurrent ketoacidosis in a teenager is rarely a problem of insulin pharmacology. Automated delivery removes some of the daily decisions. It does not remove the adolescent, the household, or the reasons the insulin was not given.

Objectives: *To review the published evidence on hybrid closed-loop insulin delivery in children and adolescents: what it achieves in glycaemic terms, what is known about its effect in those with the poorest control, what it asks of the family, and where the evidence stops.*

Material and Methods: *Narrative review of randomised and crossover trials, meta-analyses, registry analyses and psychosocial studies indexed in PubMed and the major diabetes and paediatric journals. No new patient data were generated; every estimate quoted is one published by the original investigators.*

Results: *Pooled across four randomised trials and 501 children and adolescents, of whom 323 were allocated to closed-loop therapy, time in the target range of 70 to 180 mg/dL improved by a weighted mean difference of 10.89 percentage points (95% CI 8.22 to 13.56), and participants were more likely to reach a glycosylated haemoglobin below 7 per cent (risk ratio 2.61, 95% CI 1.29 to 5.28). In very young children followed for 18 months, time in target rose 8.7 percentage points to 71 per cent with a fall in glycosylated haemoglobin of 0.4 per cent.*

Conclusion: *The glycaemic benefit is consistent, of a size that matters clinically, and reproduced across systems and age groups. What the trials do not establish is any effect on ketoacidosis, because trial populations were largely already in reasonable control and adolescents with recurrent ketoacidosis were generally not enrolled. The technology addresses the decisions; recurrent ketoacidosis usually reflects something else.*

Abbreviations: *CGM – Continuous Glucose Monitoring, DKA – Diabetic Ketoacidosis, HbA1c – Glycosylated Haemoglobin, HCL – Hybrid Closed-Loop, SAP – Sensor-Augmented Pump, TIR – Time in Range, WMD – Weighted Mean Difference.*

Keywords: *Type 1 Diabetes; Hybrid Closed-Loop; Adolescents; Diabetic Ketoacidosis; Family Support*

Introduction

Glycaemic control in type 1 diabetes deteriorates predictably during adolescence and recovers, partially, afterwards. International registry comparisons show the pattern across countries with very different baseline achievement, which suggests it is developmental rather than a failure of any particular health system [1,2]. Diabetic ketoacidosis in an established patient is the sharp end of that deterioration and is associated with insulin omission far more often than with device failure [3,4].

Hybrid closed-loop systems pair a continuous glucose sensor with an algorithm that modulates basal insulin delivery automatically, leaving meal boluses to the user. The premise is that removing most of the moment-to-moment decisions removes most of the opportunities to get them wrong [5,6].

The technology now has substantial randomised support in children and adolescents [7,8,9,10]. This review sets out what that evidence shows, what it does not address, and why an adolescent presenting with recurrent ketoacidosis sits partly outside it.

Material and Methods

Search strategy: PubMed and the archives of the major diabetes, endocrinology and paediatric journals were searched for studies of automated insulin delivery in young people. Search terms combined type 1 diabetes with hybrid closed-loop, automated insulin delivery, artificial pancreas, time in range, adolescent, ketoacidosis and adherence. Reference lists of consensus documents [11,12] were screened by hand for primary sources.

Eligibility: Studies were included if they compared automated insulin delivery with another modality in participants under 21, reported glycaemic or psychosocial outcomes with these systems, or characterised ketoacidosis and adherence in this age group. Trials in adults are cited where they establish the platform and are identified as such.

Data extraction and handling: Design, sample size, age range, comparator, baseline control and reported outcome were extracted as published. Baseline glycated haemoglobin and the trial entry criteria are reported alongside every efficacy figure, because the populations enrolled were mostly in fair control already and that bears directly on whether the results apply to a patient with recurrent ketoacidosis. No pooling or re-analysis was performed. Figures 1 to 4 and Tables 1 and 2 reproduce published values only. This is a narrative synthesis and was not registered as a systematic review.

Results

The pooled glycaemic effect: A meta-analysis of four randomised trials including 501 children and adolescents, of whom 323 were allocated to closed-loop therapy, found time in

the target range of 70 to 180 mg/dL improved by a weighted mean difference of 10.89 percentage points (95% CI 8.22 to 13.56) against control therapies (Figure 1) [13]. Roughly eleven percentage points is around two and a half hours more each day within range, which is a clinically substantial difference rather than a statistical one.




Figure 1: Improvement in time in range with hybrid closed-loop against control therapy, pooled across four randomised trials [13]

Reaching target: The same analysis found participants using closed-loop therapy more than twice as likely to achieve a glycated haemoglobin below 7 per cent (risk ratio 2.61, 95% CI 1.29 to 5.28) (Figure 2) [13]. The confidence interval is wide, reflecting the modest number of trials, but excludes no effect.




Figure 2: Likelihood of achieving glycated haemoglobin below 7 per cent with closed-loop therapy [13]

Durability and the youngest patients: Benefit is not confined to short trial periods or to older children. In very young children studied over 18 months, time in target rose by 8.7 percentage points to 71 per cent against sensor-augmented pump therapy, with a reduction in glycated haemoglobin of 0.4 per cent (Figure 3) [14,15,16]. Randomised trials of the Cambridge algorithm in participants aged 6 to 18 with baseline glycated haemoglobin around 8.2 per cent showed a lower value at six months in the closed-loop group [17], and trials of other systems in children aged 6 to 12 and in adolescents report the same direction [7,8,18], as do trials comparing activation schedules and advanced systems [33,35] and a wider meta-analysis including adults [34].




Figure 3: Time in target range over 18 months of hybrid closed-loop use in very young children [14]

The size of the evidence base: The pooled paediatric randomised evidence comprises four trials and 501 participants (Figure 4) [13]. That is a reasonable foundation for a device but a narrow one for extrapolation, and it matters most when considering populations the trials did not enrol.




Figure 4: Size of the pooled randomised evidence base in children and adolescents [13]

Study	Design and cohort	Principal reported finding
Meta-analysis, 4 RCTs [13]	501 children and adolescents, 323 on HCL	TIR +10.89 pp (8.22-13.56); HbA1c <7% RR 2.61 (1.29-5.28)
Breton et al. 2020 [7]	RCT, children aged 6-13	Increased time in range over 16 weeks against sensor-augmented pump
Ware et al. 2022 [15]	RCT, very young children	Closed-loop superior to sensor-augmented pump therapy
18-month extension [14]	Very young children, single arm	TIR 71%, +8.7 pp; HbA1c -0.4%; benefit maintained
Boughton et al. 2022 [17]	RCT, aged 6-18, baseline HbA1c 8.2%	Lower HbA1c at 6 months with closed-loop
Australian RCT 2021 [19]	172 children and adolescents, 6 months	Glycaemic and psychosocial outcomes assessed together
Bergental et al. 2021 [18]	FLAIR, adolescents and young adults	Compared two closed-loop systems directly

Table 1: Principal studies reviewed, with the finding as reported by the original investigators

What the trials did not study: Entry criteria across these studies generally required established pump use, a glycated haemoglobin within a defined band, and a family able to complete training and study procedures [7,15,17]. The Cambridge trial enrolled participants with screening values between 7.0 and 10.0 per cent [17]. An adolescent presenting with a third episode of ketoacidosis in a year typically has a value well above that range, may not be using a pump, and is often not attending clinic. Almost none of the randomised evidence describes that patient.

Why ketoacidosis happens: Recurrent ketoacidosis in an established patient is predominantly a behavioural and social phenomenon rather than a technical one. Insulin omission is the dominant precipitant, and it is associated with psychological distress, disordered relationship with food, family conflict and socioeconomic adversity [3,4,20,21]. Diabetes distress and depressive symptoms are common in adolescence and track with control [22,23]. A pump that adjusts basal delivery cannot give a bolus that was never requested, and closed-loop systems still require the user to announce meals and to keep the infusion set and sensor in place [5,24].

What technology can and cannot displace: Trials reporting psychosocial outcomes alongside glycaemic ones generally find improvements in sleep, in parental fear of hypoglycaemia and in perceived burden, which are real gains for households organised around overnight glucose checks [19,25,26]. Device discontinuation is nonetheless common in adolescence, driven by alarm fatigue, body image, the visibility of worn devices and the effort of carbohydrate counting [27,28,29]. Structured education, family-based intervention and attention to distress remain necessary alongside the hardware rather than instead of it [30,31,32].

Question	What the evidence shows	Caution
Does it improve time in range?	Yes, by about 11 percentage points	Pooled from four trials, 501 participants [13]
Does it lower HbA1c?	Yes; more reach below 7%	RR 2.61 with a wide interval [13]
Does benefit persist?	Maintained to 18 months in young children	Single-arm extension [14]
Does it prevent ketoacidosis?	Not established	Trials were not powered or populated for it [7,15,17]
Does it reduce family burden?	Reported improvements in sleep and fear	Secondary outcomes, unblinded [19,25]
Will an adolescent keep using it?	Discontinuation is common	Alarm fatigue, body image, effort [27,28]

Table 2: Practical questions about closed-loop therapy and what the current evidence supports

Discussion

The glycaemic case is settled. Automated insulin delivery improves time in range by around eleven percentage points and roughly doubles the chance of reaching a glycated haemoglobin

below 7 per cent, and it does so consistently across systems, age groups and trial designs [13,14,17,18]. For a technology, that is about as clean a result as paediatric diabetes has produced. Any child or adolescent who can access one of these systems should be offered it.

The difficulty is that the question posed by a teenager with recurrent ketoacidosis is not the question the trials answered. Those studies enrolled young people already using pumps, already within a defined glycaemic band, and supported by families able to complete study procedures [7,15,17]. Recurrent ketoacidosis in an established patient is associated with insulin omission, and omission is associated with distress, conflict and adversity rather than with the arithmetic of basal rates [3,4,20]. Extrapolating an eleven-point improvement in time in range to that patient assumes the mechanism of their poor control is the one the device addresses.

There is a reasonable counter-argument, and it should be stated. Automation reduces the number of daily actions required, and fewer required actions may mean fewer omitted ones. It also reduces the visibility of diabetes work in the household, which may lower the conflict that drives some omission. Both are plausible; neither has been demonstrated in the population that matters. What would settle it is a trial enrolling adolescents with elevated glycated haemoglobin and a history of ketoacidosis, with ketoacidosis as an endpoint. That trial has not been done, which is understandable, since it is the hardest population to recruit and retain.

The practical implication is not to withhold the technology but to be clear about what it is expected to do. Offering a closed-loop system to an adolescent whose insulin is not being given, without addressing why it is not being given, risks two failures: the ketoacidosis recurs, and the family concludes that the best available technology did not work for them. The psychosocial and family-directed interventions in this literature are less impressive individually than the device trials, and are more likely to be the active ingredient in this specific presentation [30,31,32].

Study limitations

The limitations belong to the underlying literature. The pooled paediatric evidence is four randomised trials and 501 participants, with heterogeneity in systems, comparators and duration [13]. Entry criteria excluded the poorest control and required existing pump use, so the trials are not representative of the population with recurrent ketoacidosis [7,15,17]. Blinding is impossible, and psychosocial outcomes are particularly vulnerable to that [19,25]. Follow-up rarely exceeds six months, with one 18-month single-arm extension [14]. Ketoacidosis is too infrequent for these sample sizes to assess, so its absence from the results is uninformative rather than reassuring. Trial participants receive intensive support that routine care does not provide, and real-world discontinuation rates are higher than trial

completion rates suggest [27,28]. And the evidence comes almost entirely from high-income settings with reimbursed device access [1,2].

Conclusion

Hybrid closed-loop delivery is the most effective glycaemic intervention available to children and adolescents with type 1 diabetes, improving time in range by roughly eleven percentage points and doubling the likelihood of reaching target. That case is made. What has not been shown is that it prevents recurrent ketoacidosis in adolescents, and the reason is that the trials enrolled young people who were already doing reasonably well. For a teenager admitted for the third time in a year, the technology is worth offering and is unlikely to be sufficient on its own, because the insulin that was not delivered was usually not delivered for reasons an algorithm cannot reach.

Acknowledgements

The authors thank the investigators whose trials are drawn together in this article, and the families who tolerated masked continuous glucose monitoring and repeated study visits through periods of adolescence.

Funding

No dedicated funding from public, commercial or charitable sources supported the preparation of this review.

Conflict of Interest

The authors declare no competing interests.

Ethics Statement

This article reviews previously published studies only. No new patient data were collected and no participants were recruited, so ethical approval was not required.

References

1. Anderzen J, Hermann JM, Samuelsson U, Charalampopoulos D, Svensson J, Skrivarhaug T, et al. "International benchmarking in type 1 diabetes: large difference in childhood HbA1c between eight high-income countries but similar rise during adolescence". *Pediatr Diabetes* 21(2020):621-627.
2. Foster NC, Beck RW, Miller KM, Clements MA, Rickels MR, DiMeglio LA, et al. "State of type 1 diabetes management and outcomes from the T1D Exchange in 2016-2018". *Diabetes Technol Ther* 21(2019):66-72.
3. Cengiz E, Xing D, Wong JC, Wolfsdorf JJ, Haymond MW, Rewers A, et al. "Severe hypoglycemia and diabetic ketoacidosis among youth with type 1 diabetes in the T1D Exchange clinic registry". *Pediatr Diabetes* 14(2013):447-454.

4. Morris AD, Boyle DIR, McMahon AD, Greene SA, MacDonald TM, Newton RW. "Adherence to insulin treatment, glycaemic control, and ketoacidosis in insulin-dependent diabetes mellitus". *Lancet* 350(1997):1505-1510.
5. Boughton CK, Hovorka R. "Advances in artificial pancreas systems". *Sci Transl Med* 11(2019):eaaw4949.
6. Sherr JL. "Closing the loop on managing youth with type 1 diabetes: children are not just small adults". *Diabetes Care* 41(2018):1572-1578.
7. Breton MD, Kanapka LG, Beck RW, Ekhlaspour L, Forlenza GP, Cengiz E, et al. "A randomized trial of closed-loop control in children with type 1 diabetes". *N Engl J Med* 383(2020):836-845.
8. Brown SA, Kovatchev BP, Raghinaru D, Lum JW, Buckingham BA, Kudva YC, et al. "Six-month randomized, multicenter trial of closed-loop control in type 1 diabetes". *N Engl J Med* 381(2019):1707-1717.
9. Wadwa RP, Reed ZW, Buckingham BA, DeBoer MD, Ekhlaspour L, Forlenza GP, et al. "Trial of hybrid closed-loop control in young children with type 1 diabetes". *N Engl J Med* 388(2023):991-1001.
10. Tauschmann M, Thabit H, Bally L, Allen JM, Hartnell S, Wilinska ME, et al. "Closed-loop insulin delivery in suboptimally controlled type 1 diabetes: a multicentre, 12-week randomised trial". *Lancet* 392(2018):1321-1329.
11. Battelino T, Danne T, Bergenstal RM, Amiel SA, Beck R, Biester T, et al. "Clinical targets for continuous glucose monitoring data interpretation: recommendations from the international consensus on time in range". *Diabetes Care* 42(2019):1593-1603.
12. Sherr JL, Schoelwer M, Dos Santos TJ, Reddy L, Biester T, Galderisi A, et al. "ISPAD clinical practice consensus guidelines 2022: diabetes technologies: insulin delivery". *Pediatr Diabetes* 23(2022):1406-1431.
13. Meta-analysis Group. "Efficacy of the hybrid closed-loop insulin delivery system in children and adolescents with type 1 diabetes: a meta-analysis with trial sequential analysis". *Arch Endocrinol Metab* (2024).
14. Ware J, Boughton CK, Allen JM, Wilinska ME, Tauschmann M, Denvir L, et al. "Eighteen-month hybrid closed-loop use in very young children with type 1 diabetes: a single-arm multicenter trial". *Diabetes Care* 47(2024):2189-2193.
15. Ware J, Allen JM, Boughton CK, Wilinska ME, Hartnell S, Thankamony A, et al. "Randomized trial of closed-loop control in very young children with type 1 diabetes". *N Engl J Med* 386(2022):209-219.
16. Tauschmann M, Allen JM, Nagl K, Fritsch M, Yong J, Metcalfe E, et al. "Home use of day-and-night hybrid closed-loop insulin delivery in very young children: a multicenter, 3-week, randomized trial". *Diabetes Care* 42(2019):594-600.
17. Boughton CK, Hartnell S, Thabit H, Poettler T, Herzig D, Wilinska ME, et al. "Cambridge hybrid closed-loop algorithm in children and adolescents with type 1 diabetes: a multicentre 6-month randomised controlled trial". *Lancet Digit Health* 4(2022):e245-e255.
18. Bergenstal RM, Nimri R, Beck RW, Criego A, Laffel L, Schatz D, et al. "A comparison of two hybrid closed-loop systems in adolescents and young adults with type 1 diabetes (FLAIR): a multicentre, randomised, crossover trial". *Lancet* 397(2021):208-219.
19. Abraham MB, de Bock M, Smith GJ, Dart J, Fairchild JM, King BR, et al. "Effect of a hybrid closed-loop system on glycemic and psychosocial outcomes in children and adolescents with type 1 diabetes: a randomized clinical trial". *JAMA Pediatr* 175(2021):1227-1235.
20. Garrett CJ, Choudhary P, Amiel SA, Fonagy P, Ismail K. "Recurrent diabetic ketoacidosis and a brief history of brittle diabetes research: contemporary and past evidence in diabetic ketoacidosis research including mortality, mental health and prevention". *Diabet Med* 36(2019):1329-1335.
21. Rewers A, Chase HP, Mackenzie T, Walravens P, Roback M, Rewers M, et al. "Predictors of acute complications in children with type 1 diabetes". *JAMA* 287(2002):2511-2518.
22. Hagger V, Hendrieckx C, Sturt J, Skinner TC, Speight J. "Diabetes distress among adolescents with type 1 diabetes: a systematic review". *Curr Diab Rep* 16(2016):9.
23. Buchberger B, Huppertz H, Krabbe L, Lux B, Mattivi JT, Siafarikas A. "Symptoms of depression and anxiety in youth with type 1 diabetes: a systematic review and meta-analysis". *Psychoneuroendocrinology* 70(2016):70-84.
24. Kanapka LG, Wadwa RP, Breton MD, Ruedy KJ, Ekhlaspour L, Forlenza GP, et al. "Extended use of the Control-IQ closed-loop control system in children with type 1 diabetes". *Diabetes Care* 44(2021):473-478.
25. Van Name MA, Hilliard ME, Boyle CT, Miller KM, DeSalvo DJ, Anderson BJ, et al. "Nighttime is the worst time: parental fear of hypoglycemia in young children with type 1 diabetes". *Pediatr Diabetes* 19(2018):114-120.
26. Barnard KD, Wysocki T, Ully V, Mader JK, Pieber TR, Thabit H, et al. "Closing the loop in adults, children and adolescents with suboptimally controlled type 1 diabetes under free living conditions: a psychosocial substudy". *J Diabetes Sci Technol* 11(2017):1080-1088.

27. Wong JC, Foster NC, Maahs DM, Raghinaru D, Bergenstal RM, Ahmann AJ, et al. "Real-time continuous glucose monitoring among participants in the T1D Exchange clinic registry". *Diabetes Care* 37(2014):2702-2709.
28. Messer LH, Tanenbaum ML, Cook PF, Wong JJ, Hanes SJ, Driscoll KA, et al. "Cost, hassle, and on-body experience: barriers to diabetes device use in adolescents and potential intervention targets". *Diabetes Technol Ther* 22(2020):760-767.
29. Tanenbaum ML, Hanes SJ, Miller KM, Naranjo D, Bensen R, Hood KK. "Diabetes device use in adults with type 1 diabetes: barriers to uptake and potential intervention targets". *Diabetes Care* 40(2017):181-187.
30. Hood KK, Rohan JM, Peterson CM, Drotar D. "Interventions with adherence-promoting components in pediatric type 1 diabetes: meta-analysis of their impact on glycemic control". *Diabetes Care* 33(2010):1658-1664.
31. Wysocki T, Harris MA, Buckloh LM, Mertlich D, Lochrie AS, Taylor A, et al. "Randomized trial of behavioral family systems therapy for diabetes: maintenance of effects on diabetes outcomes in adolescents". *Diabetes Care* 30(2007):555-560.
32. Katz ML, Volkening LK, Butler DA, Anderson BJ, Laffel LM. "Family-based psychoeducation and care ambassador intervention to improve glycemic control in youth with type 1 diabetes: a randomized trial". *Pediatr Diabetes* 15(2014):142-150.
33. Renard E, Tubiana-Rufi N, Bonnemaïson-Gilbert E, Coutant R, Dalla-Vale F, Farret A, et al. "Outcomes of hybrid closed-loop insulin delivery activated 24/7 versus evening and night in free-living prepubertal children with type 1 diabetes: a multicentre, randomized clinical trial". *Diabetes Obes Metab* 24(2022):511-521.
34. Jiao X, Shen Y, Chen Y. "Better TIR, HbA1c, and less hypoglycemia in closed-loop insulin system in patients with type 1 diabetes: a meta-analysis". *BMJ Open Diabetes Res Care* 10(2022):e002633.
35. Collyns OJ, Meier RA, Betts ZL, Chan DSH, Frampton C, Frewen CM, et al. "Improved glycemic outcomes with Medtronic MiniMed advanced hybrid closed-loop delivery: results from a randomized crossover trial comparing automated insulin delivery with predictive low glucose suspend in people with type 1 diabetes". *Diabetes Care* 44(2021):969-975.

Citation: Zielinska K, Bauer L, Nieminen E. "Hybrid Closed-Loop Insulin Delivery in an Adolescent with Recurrent Diabetic Ketoacidosis: A Narrative Review of Technology, Adherence and Family Support." *J Clin Case Stud* (2024): 001-006 doi: 10.5281/zenodo.22236494