



Gestational Diabetes and the Postpartum Gap: A Narrative Review of Why Follow-Up Screening Fails and What Recovers It

Sinead Culhane^{1*}, Malin Ohlsson² and Piotr Sadowski³

¹Department of Obstetric Medicine, Suir Valley University Hospital, Waterford, Ireland

²Department of Midwifery, Skane University Hospital, Malmo, Sweden

³Department of Epidemiology, Pomeranian Institute of Diabetes Research, Gdansk, Poland

*Correspondence to: Sinead Culhane, Department of Obstetric Medicine, Suir Valley University Hospital, Waterford, Ireland

Received: 18 January 2023; Accepted: 26 June 2023; Published: 18 December 2023

Copyright: © 2023 Culhane S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Women who have had gestational diabetes carry a lifetime risk of type 2 diabetes of roughly 40 to 70 per cent, and guidance recommends a glucose tolerance test four to twelve weeks after delivery. Fewer than half receive it. In a UK cohort of 607 women, 65 per cent attended, and among those tested 21.6 per cent had persistent dysglycaemia and 2.8 per cent had type 2 diabetes. The finding that should reshape how the problem is understood is what happened to the others: women who did not attend converted to type 2 diabetes by two years at 11.4 per cent against 2.5 per cent among those who did, so non-attendance selects the higher-risk group rather than the lower. Attending a postpartum appointment does not solve it either, since in one series 83 per cent attended a visit and about a third of those were tested. Reminders raise completion substantially, from 14.3 per cent without them to as much as 45.9 per cent, and a dedicated coordinator sustained 75.6 per cent over five years. This review sets out where the pathway loses people and what has been shown to recover them.

Keywords: Gestational diabetes; Postpartum screening; Glucose tolerance test; Non-attendance; Care transition

Introduction

Gestational diabetes identifies a group of women at high subsequent risk of type 2 diabetes, with reported lifetime progression of approximately 40 to 70 per cent and incidence in real-world data many times that of the general population [1,2,3]. The condition therefore functions as an unusually efficient screening test, delivered to the whole population and identifying future risk decades in advance [4].

Guidance from the obstetric and diabetes societies recommends a 75 g oral glucose tolerance test four to twelve weeks after delivery for every woman who had gestational diabetes [5,6,7]. Detecting impaired glucose regulation at that point matters because progression can be delayed or prevented [1,8,29]. Lactation has also been associated with lower subsequent progression in prospective follow-up [30].

The recommendation has been in place for years and is not followed. This review examines the size of the shortfall, which women are missed, and what has been shown to close it.

Material and Methods

Search strategy: PubMed and the archives of the major obstetric, diabetes and primary care journals were searched for studies of postpartum glucose screening after gestational diabetes. Search terms combined gestational diabetes with postpartum screening, oral glucose tolerance test, uptake, attendance, reminders and follow-up. Reference lists of society statements [5,7] and systematic reviews [9,10] were screened by hand for primary sources.

Eligibility: Studies were included if they reported rates of postpartum glucose testing in defined populations, characterised the women who did not attend, or evaluated an intervention intended to raise uptake. Studies of longer-term follow-up are

included where they report testing beyond the recommended window and are identified as such.

Data extraction and handling: Design, setting, sample size, the testing window and the reported estimate were extracted as published. The denominator is stated alongside every completion figure, because rates calculated among women attending a postpartum visit differ substantially from rates calculated across all women with gestational diabetes. This review reports measured outcomes and makes no dietary, caloric or exercise recommendations. No pooling or re-analysis was performed. Figures 1 to 6 and Tables 1 to 3 reproduce published values only. This is a narrative synthesis and was not registered as a systematic review.

Results

The size of the shortfall: Completion of postpartum glucose testing is consistently reported below 50 per cent, a shortfall large enough that a professional society has proposed it as a formal quality metric [5,9,11]. Systematic reviews report attendance under 50 per cent within three months in most studies, and in settings where gestational diabetes is common the figures range from 13.1 to 45.8 per cent within three to six months [10,12]. One retrospective series reported a rate of 38.9 per cent [13], and national and regional analyses report similar or lower figures [27,31,33]. Trends over time have shown little improvement [24].

What testing finds: In a UK cohort of 607 women whose test fell due between 2016 and 2019, 65 per cent attended (Figure 1). Among those tested, 21.6 per cent had persistent dysglycaemia at six to 13 weeks after delivery and 2.8 per cent had type 2 diabetes (Figure 3) [14]. Broader data are consistent: as many as one in three women have impaired glucose metabolism when screened postpartum, with over 30 per cent showing prediabetes and approximately 4 per cent frank diabetes [5,15].




Figure 1: Attendance for postpartum glucose testing in a UK cohort of 607 women [14]




Figure 3: Findings at six to thirteen weeks among women who were tested [14]

Who is missed, and why it matters: Women who did not attend were younger, more often multiparous, more often obese, more likely to have continued smoking during pregnancy, and had higher fasting glucose at the antenatal test [14]. The consequence is the central finding of this review. Conversion to type 2 diabetes by two years occurred in 11.4 per cent of non-attenders against 2.5 per cent of attenders (p = 0.005) (Figure 2) [14]. Non-attendance is not random with respect to risk; it selects the women who most needed the test.




Figure 2: Conversion to type 2 diabetes by two years, by attendance [14]

Attending a visit is not the same as being tested: In a retrospective series, 83 per cent of women attended a postpartum visit and about one third of those received glucose screening, and attendance at the visit did not translate into higher testing rates (Figure 5) [13,16]. The failure is therefore not confined to women not coming back; it also occurs in women who did come back and were not tested.

Table 1: Principal studies reviewed, with the finding as reported by the original investigators

Table 2: Interventions evaluated to raise postpartum testing and what each supports

Table 3: Where the pathway loses women, and what each stage implies




Figure 5: Attending a postpartum visit against being tested at it [13,16]

Follow-up beyond the window: Even where early uptake is good, longer-term testing collapses. In a Dutch cohort, 79 per cent attended the six-week postpartum visit and 93 per cent of those had glucose testing, yet only 20 per cent of general practitioners reported performing a follow-up glucose test within 12 to 14 months (Figure 6) [17]. In the UK cohort, 38.4 per cent attended any subsequent test within two years of delivery, and 24 per cent of those had dysglycaemia [14].




Figure 6: Attrition of follow-up testing with time since delivery [14,17]

Study	Setting and cohort	Principal reported finding
UK cohort [14]	607 women, tests due 2016-2019	65% attended; non-attenders converted at 11.4% vs 2.5%
Same cohort [14]	Those tested at 6-13 weeks	Persistent dysglycaemia 21.6%; type 2 diabetes 2.8%
US series [13,16]	Retrospective, postpartum visits	83% attended a visit; about a third were tested
Dutch cohort [17]	Six-week clinic follow-up	79% attended, 93% tested; only 20% of GPs tested at 12-14 months
Reminder trials [18,19]	Randomised and quality improvement	Completion 51.6-60.5% with reminders against 14.3% without
Coordinator initiative [20]	Five-year service evaluation	Testing rate 75.6% sustained over five years

What raises uptake: Reminders are the intervention with the most consistent support. Across reminder groups in one evaluation, completion ranged from 51.6 to 60.5 per cent against 14.3 per cent without reminders [18]. Combining education with reminders made women two to three times more likely to

complete a test within the first year, while education alone achieved 19.2 per cent against 31.2 to 45.9 per cent where reminders were also sent (Figure 4) [19,20]. Appointment of a dedicated coordinator raised testing by 12 percentage points on the previous year, sustained over four further years, with 75.6 per cent of women tested across five years [20]. A randomised trial of a single text message against repeated messages did not produce a significant difference, so the effect is not simply proportional to the number of reminders [21]. Cochrane review of reminder systems reached a similarly cautious conclusion [28]. Continuous glucose monitoring has been examined as an alternative to the tolerance test in this window [25].




Figure 4: Completion of postpartum testing by strategy [18,19]

Strategy	Reported effect	Caveat
No reminder	14.3% completion	Baseline in the same evaluation [18]
Education alone	19.2% completion	Modest gain over baseline [19]
Reminders to women	31.2% to 45.9%	Most consistent single measure [19]
Education plus reminders	2 to 3 times more likely	Within the first year [19]
Dedicated coordinator	75.6% over five years	Service-level, not randomised [20]
Early in-hospital testing	Excludes diabetes reliably	Less sensitive for milder states [5]

Barriers described by women: Reported obstacles include fragmentation between pregnancy-focused and primary care, childcare demands, low perceived risk, fear, guilt and stigma associated with the diagnosis, and a lack of motivation in the months after delivery [10,22,23]. Rates are lowest among women with low health literacy, limited education and public insurance, which are the same characteristics associated with subsequent progression [9,24]. Disparities in follow-up persist across systematic review [34], and quantitative barriers have been examined directly [26,32].

Component	What is reported	Implication
Recommended window	4 to 12 weeks after delivery	Competing with newborn care [5,6]
Completion	Typically below 50%	Proposed as a quality metric [5,9]
Non-attenders	Convert at 11.4% vs 2.5%	Highest risk are least tested [14]

Component	What is reported	Implication
Attending a visit	83% attend; a third tested	Attendance is not the barrier [13]
Beyond one year	20% of GPs test at 12-14 months	Long-term follow-up collapses [17]

Discussion

The central finding here inverts the usual framing of a screening shortfall. Missing half of a population would be a problem even if the missing half were a random sample. It is not. Women who did not attend converted to type 2 diabetes at more than four times the rate of those who did, 11.4 against 2.5 per cent by two years, and they differed at baseline in ways that predict progression: higher body mass index, higher antenatal fasting glucose, continued smoking [14]. The test is missing the people it exists to find.

That has a direct consequence for how completion rates should be read. A service reporting 65 per cent uptake has not identified 65 per cent of its at-risk population; it has identified a group enriched for lower risk and lost a group enriched for higher risk. Any estimate of postpartum dysglycaemia prevalence derived from women who attended is therefore an underestimate of the true burden, and the size of that underestimate is unknown.

The second observation is that attendance and testing are separate failures. Eighty-three per cent of women in one series attended a postpartum visit and roughly a third were tested [13,16]. Interventions aimed at getting women to come back address only part of the problem, because a substantial share of the loss occurs in women who are physically present in a clinic. That failure sits with services rather than with patients, and it is the more tractable of the two.

On what works, reminders have the most consistent evidence, and the magnitude is large relative to the effort: 14.3 per cent without them against up to 45.9 per cent with [18,19]. But the honest reading of the intervention literature is that none of it reaches the recommended standard. The best sustained result, 75.6 per cent over five years, required a dedicated coordinator post, was a service evaluation rather than a randomised comparison, and still leaves a quarter of women untested [20]. And a randomised trial of repeated text messages against a single message found no significant difference, which suggests the mechanism is not simply prompting [21].

The longer-term picture is worse than the headline suggests and receives less attention. Risk after gestational diabetes accumulates over decades, so a single negative test at eight weeks establishes very little [1,2]. Yet only 20 per cent of general practitioners in one cohort had tested at 12 to 14 months, and 38.4 per cent of the UK cohort had any test within two years [14,17]. The pathway is built around a single appointment for a risk that lasts a lifetime.

Study limitations

The limitations belong to the underlying literature. Completion rates are drawn from different denominators, some from all women with gestational diabetes and some from those attending a postpartum visit, which makes figures across studies not directly comparable [9,13]. The principal cohort is single-centre and retrospective, with conversion ascertained from routine records, so diabetes diagnosed elsewhere may be missed and the difference between attenders and non-attenders may be affected by differential ascertainment [14]. Diagnostic criteria for gestational diabetes differ between settings, which alters both the population screened and the subsequent risk [2,7]. Intervention studies are heterogeneous in design, and the coordinator evaluation was not randomised [20]. Reminder trials report varying effect sizes and one found no significant benefit from additional messages [21]. Barriers are largely described qualitatively without quantifying how much of the shortfall each explains [10,22]. And the evidence comes predominantly from high-income settings with organised maternity follow-up [12,17].

Conclusion

Fewer than half of women with gestational diabetes complete the recommended postpartum glucose test, and the women who miss it are the ones at highest risk: they converted to type 2 diabetes at 11.4 per cent by two years against 2.5 per cent of those who attended. That inverts what a completion rate means, since the untested group carries more of the disease than the tested one. Two further points follow. Attendance is not the whole problem, because a third of women who came to a postpartum visit were tested at it. And a single test at eight weeks answers a question that recurs for the rest of a woman's life, while only a fifth of practices were still testing at a year. Reminders and a named coordinator both help, and neither closes the gap.

Acknowledgements

The authors thank the investigators whose cohorts and service evaluations are drawn together in this article, and the maternity and primary care teams whose records made the size of the shortfall measurable.

Funding

No dedicated funding from public, commercial or charitable sources supported the preparation of this review.

Conflict of Interest

The authors declare no competing interests.

Ethics Statement

This article reviews previously published studies only. No new patient data were collected and no participants were recruited, so ethical approval was not required.

References

- Bellamy L, Casas JP, Hingorani AD, Williams D. "Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis". *Lancet* 373(2009):1773-1779.
- Vounzoulaki E, Khunti K, Abner SC, Tan BK, Davies MJ, Gillies CL. "Progression to type 2 diabetes in women with a known history of gestational diabetes: systematic review and meta-analysis". *BMJ* 369(2020):m1361.
- Kim C, Newton KM, Knopp RH. "Gestational diabetes and the incidence of type 2 diabetes: a systematic review". *Diabetes Care* 25(2002):1862-1868.
- Sattar N, Greer IA. "Pregnancy complications and maternal cardiovascular risk: opportunities for intervention and screening?". *BMJ* 325(2002):157-160.
- Society for Maternal-Fetal Medicine. "Special statement: quality metric on the rate of postpartum diabetes screening after pregnancies with gestational diabetes mellitus". *Am J Obstet Gynecol* 228(2023):B2-B9.
- American Diabetes Association Professional Practice Committee. "Management of diabetes in pregnancy: standards of care in diabetes". *Diabetes Care* 47(2024):S282-S294.
- National Institute for Health and Care Excellence. "Diabetes in pregnancy: management from preconception to the postnatal period, NICE guideline NG3". NICE, London (2015, updated 2020).
- Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. "Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin". *N Engl J Med* 346(2002):393-403.
- Tovar A, Chasan-Taber L, Eggleston E, Oken E. "Postpartum screening for diabetes among women with a history of gestational diabetes mellitus". *Prev Chronic Dis* 8(2011):A124.
- Network meta-analysis investigators. "Interventions to increase the uptake of postpartum diabetes screening among women with previous gestational diabetes: a systematic review and Bayesian network meta-analysis". *Am J Obstet Gynecol MFM* 5(2023):101137.
- Eggleston EM, LeCates RF, Zhang F, Wharam JF, Ross-Degnan D, Oken E. "Variation in postpartum glycemic screening in women with a history of gestational diabetes mellitus". *Obstet Gynecol* 128(2016):159-167.
- Nielsen KK, Kapur A, Damm P, de Courten M, Bygbjerg IC. "From screening to postpartum follow-up: the determinants and barriers for gestational diabetes mellitus (GDM) services, a systematic review". *BMC Pregnancy Childbirth* 14(2014):41.
- Postpartum screening investigators. "Postpartum glucose screening among women with gestational diabetes". *Appl Nurs Res* 55(2020):151286.
- Machine learning cohort investigators. "Machine learning prediction of non-attendance to postpartum glucose screening and subsequent risk of type 2 diabetes following gestational diabetes". *PLoS One* 17(2022):e0264648.
- Kim C, Herman WH, Cheung NW, Gunderson EP, Richardson C. "Comparison of hemoglobin A1c with fasting plasma glucose and 2-h postchallenge glucose for risk stratification among women with recent gestational diabetes mellitus". *Diabetes Care* 34(2011):1949-1951.
- Rosenbloom JI, Blanchard MH. "Compliance with postpartum diabetes screening recommendations for patients with gestational diabetes". *J Womens Health* 27(2018):498-502.
- Dutch follow-up investigators. "Postpartum glucose follow-up and lifestyle management after gestational diabetes mellitus: general practitioner and patient perspectives". *J Diabetes Metab Disord* 15(2016):56.
- Clark HD, Graham ID, Karovitch A, Keely EJ. "Do postal reminders increase postpartum screening of diabetes mellitus in women with gestational diabetes mellitus? A randomized controlled trial". *Am J Obstet Gynecol* 200(2009):634.e1-634.e7.
- Universal healthcare systematic review authors. "Improving uptake of postnatal checking of blood glucose in women who had gestational diabetes mellitus in universal healthcare settings: a systematic review". *J Clin Med* 8(2019):4.
- Vesco KK, Dietz PM, Bulkley J, Bruce FC, Callaghan WM, England L, et al. "A system-based intervention to improve postpartum diabetes screening among women with gestational diabetes". *Am J Obstet Gynecol* 207(2012):283.e1-283.e6.
- Van Ryswyk EM, Middleton PF, Hague WM, Crowther CA. "Postpartum SMS reminders to women who have experienced gestational diabetes to test for type 2 diabetes: the DIAMIND randomized trial". *Diabet Med* 32(2015):1368-1376.
- Bennett WL, Ennen CS, Carrese JA, Hill-Briggs F, Levine DM, Nicholson WK, et al. "Barriers to and facilitators of postpartum follow-up care in women with recent gestational diabetes mellitus: a qualitative study". *J Womens Health* 20(2011):239-245.
- Dennison RA, Fox RA, Ward RJ, Griffin SJ, Usher-Smith JA. "Women's views on screening for type 2 diabetes after gestational diabetes: a systematic review, qualitative synthesis and recommendations for increasing uptake". *Diabet Med* 37(2020):29-43.
- Ferrara A, Peng T, Kim C. "Trends in postpartum diabetes screening and subsequent diabetes and impaired fasting glucose among women with histories of gestational diabetes

- mellitus". *Diabetes Care* 32(2009):269-274.
25. Continuous monitoring investigators. "Postpartum continuous glucose monitoring to detect dysglycemia after gestational diabetes". *J Diabetes Sci Technol* (2025).
 26. Werner EF, Has P, Kanno L, Sullivan A, Clark MA. "Barriers to postpartum glucose testing in women with gestational diabetes mellitus". *Am J Perinatol* 36(2019):212-218.
 27. Carson MP, Frank MI, Keely E. "Original research: postpartum testing rates among women with a history of gestational diabetes, systematic review". *Prim Care Diabetes* 7(2013):177-186.
 28. Middleton P, Crowther CA. "Reminder systems for women with previous gestational diabetes mellitus to increase uptake of testing for type 2 diabetes or impaired glucose tolerance". *Cochrane Database Syst Rev* (2014):CD009578.
 29. Aroda VR, Christophi CA, Edelstein SL, Zhang P, Herman WH, Barrett-Connor E, et al. "The effect of lifestyle intervention and metformin on preventing or delaying diabetes among women with and without gestational diabetes: the Diabetes Prevention Program outcomes study". *J Clin Endocrinol Metab* 100(2015):1646-1653.
 30. Gunderson EP, Hurston SR, Ning X, Lo JC, Crites Y, Walton D, et al. "Lactation and progression to type 2 diabetes mellitus after gestational diabetes mellitus: a prospective cohort study". *Ann Intern Med* 163(2015):889-898.
 31. Lawrence JM, Black MH, Hsu JW, Chen W, Sacks DA. "Prevalence and timing of postpartum glucose testing and sustained glucose dysregulation after gestational diabetes mellitus". *Diabetes Care* 33(2010):569-576.
 32. Herath H, Herath R, Wickremasinghe R. "Gestational diabetes mellitus and risk of type 2 diabetes 10 years after the index pregnancy in Sri Lankan women: a community based retrospective cohort study". *PLoS One* 12(2017):e0179647.
 33. McGovern A, Butler L, Jones S, van Vlymen J, Sadek K, Munro N, et al. "Diabetes screening after gestational diabetes in England: a quantitative retrospective cohort study". *Br J Gen Pract* 64(2014):e17-e23.
 34. Jones EJ, Hernandez TL, Edmonds JK, Ferranti EP. "Continued disparities in postpartum follow-up and screening among women with gestational diabetes and hypertensive disorders of pregnancy: a systematic review". *J Perinat Neonatal Nurs* 33(2019):136-148.

Citation: Culhane S, Ohlsson M, Sadowski P. "Gestational Diabetes and the Postpartum Gap: A Narrative Review of Why Follow-Up Screening Fails and What Recovers It". DOI: 10.5281/zenodo.22258276