



Psychosocial Empowerment Programs and Their Effect on Postpartum Mental Health

Fernandez M^{1*}, Salgado P² and Valdes R³

¹Department of Clinical Research, Seville Institute of Gynecological Research, Portugal Scopus Author ID: 57510843301 Email: mateo.fernandez@seville.edu
ORCID: <https://orcid.org/3747-5350-2791-8741>

²Department of Translational Medicine, Porto Institute of Maternal Medicine, Portugal Scopus Author ID: 57956147218
Email: pilar.salgado@porto.edu
ORCID: <https://orcid.org/1865-6839-1072-1607>

³Department of Clinical Research, Seville Institute of Gynecological Research, Spain Scopus Author ID: 57754105645
Email: rodrigo.valdes@seville.edu
ORCID: <https://orcid.org/0666-3747-9855-310X>

***Correspondence to:** Mateo Fernandez, Department of Clinical Research, Seville Institute of Gynecological Research, Portugal Scopus Author ID: 57510843301 Email: mateo.fernandez@seville.edu

Received: 8 January 2026; **Accepted:** 25 March 2026; **Published:** 17 June 2026

Copyright: © 2026 MF, et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Postpartum depression affects a substantial share of new mothers worldwide, and the strongest predictors identified in the epidemiological literature -- poor social support, exposure to intimate partner violence, and unplanned pregnancy -- are psychosocial rather than purely biological, motivating a class of interventions built around psychosocial empowerment: peer support, community health worker-delivered structured therapy, group-based prenatal and postnatal care, and targeted self-management programs designed to increase a mother's perceived control, self-efficacy, and social connectedness. This article synthesizes that literature, grounding it in Zimmerman's psychological empowerment framework and Bandura's self-efficacy theory, and tracing the evidence from the Thinking Healthy Programme's origins as a single rural Pakistani trial through its cross-cultural adaptation into routine antenatal care in Turkey and China, alongside independently convergent meta-analyses of peer support, a growing group prenatal care trial literature, and emerging evidence for compounded-risk populations facing intimate partner violence. Read together, the evidence supports psychosocial empowerment programs as a genuinely effective, moderately-sized, and increasingly portable intervention class for postpartum mental health, while leaving direct mechanistic testing of the empowerment pathway itself, dose-response and component-isolation questions, and the field's limited engagement with the structural conditions underlying psychosocial risk as an open agenda for future work.

Keywords: postpartum depression; psychosocial empowerment; peer support; maternal mental health; group prenatal care

1. Introduction

Postpartum depression is, by most global estimates, one of the most common complications of childbearing, and yet it remains one of the most inconsistently prevented and treated. A woman who develops gestational diabetes or preeclampsia enters a well-worn clinical pathway with defined screening thresholds, referral triggers, and follow-up protocols. A woman who develops postpartum depression, by contrast, often encounters a fragmented patchwork of screening tools that may or may not be administered, brief pharmacological consultations that may or may not be accessible, and a broader silence around the social conditions — isolation, loss of autonomy, unequal division of infant care labor, exposure to violence, financial precarity — that shape whether depressive symptoms take hold and how long they persist. Over the past two decades, a substantial and increasingly rigorous body of intervention research has tried to close that gap not through pharmacology but through psychosocial empowerment: structured programs that aim to increase a new mother's sense of control, self-efficacy, social connectedness, and decision-making capacity around her own health and her infant's care, on the theory that depressive symptoms in the postpartum period are not simply a biochemical event happening to a passive patient but are substantially shaped by, and responsive to, her social and psychological position.

This article synthesizes that literature. We are concerned specifically with psychosocial empowerment programs — peer support models, community health worker-delivered psychotherapy, group-based antenatal and postnatal care, and structured self-management or psychoeducation programs explicitly designed around increasing agency and coping capacity — and with what the accumulated trial and meta-analytic evidence says about their effect on postpartum mental health outcomes, principally depressive and anxiety symptoms measured via validated screening instruments such as the Edinburgh Postnatal Depression Scale [1]. We do not cover pharmacological treatment of postpartum depression, nor do we cover severe postpartum psychosis, both of which fall outside the psychosocial-empowerment frame and require different literatures entirely.

The case for taking empowerment seriously as a mechanism, rather than treating it as a vague wellness

buzzword, rests on two converging bodies of theory. The first is Zimmerman's psychological empowerment framework, which distinguishes intrapersonal empowerment (perceived control, self-efficacy, motivation to exert influence over one's own life) from interactional empowerment (the skills and critical awareness needed to actually navigate one's environment) and behavioral empowerment (the actions a person takes as a result) [2]. The second is Bandura's social cognitive account of health promotion, which places self-efficacy — a person's belief in their capacity to execute the behaviors needed to produce a given outcome — at the center of virtually every successful health behavior change effort, postpartum mental health included [3]. Put together, these frameworks predict something specific and testable: that an intervention which measurably increases a new mother's perceived control, coping skill, and social support should, if the theory holds, produce a measurable downstream reduction in depressive and anxiety symptomatology. The question this article asks is how well that prediction has actually held up against two decades of trial data.

We organize the discussion around five themes: the epidemiological scope of postpartum mental health difficulty and the specific psychosocial risk factors that motivate an empowerment-based response; the theoretical architecture of empowerment as applied to maternal mental health; the peer-support and community health worker literature, which is by far the most mature and most rigorously trialed strand of this work; group prenatal and postnatal care models, which operationalize empowerment structurally rather than individually; and programs targeting women facing compounded risk, particularly intimate partner violence, where empowerment language carries a more explicit political and clinical weight. We close with an honest accounting of where the evidence is strong, where it is suggestive but thin, and where the field still owes practitioners better answers than it currently has.

2. Postpartum Mental Health: Scope and the Case for Psychosocial Approaches

Estimates of postpartum depression prevalence vary considerably by population, screening instrument, and timing of assessment, but the consistent finding across dozens of systematic reviews is that it is common, not rare. A 2021 systematic review and meta-analysis of

eleven Ethiopian studies covering 7,582 women estimated a pooled postpartum depression prevalence of 22.08% (95% CI 17.66-26.49%), notably higher than commonly cited high-income-country estimates in the 10-15% range, and identified three factors carrying particularly strong statistical association with depression: poor social support (odds ratio 3.57), exposure to domestic violence (odds ratio 3.14), and an unplanned pregnancy (odds ratio 2.84) [4]. A separate 2025 meta-analysis synthesizing 122 studies and more than 560,000 women across 43 countries found that roughly one in ten women experience comorbid anxiety and depression during pregnancy and the postpartum period specifically, underscoring that postpartum mental health difficulty is rarely a single, isolated diagnostic event but frequently a co-occurring cluster of mood and anxiety symptoms [5]. A parallel global synthesis focused specifically on rural populations similarly found elevated and heterogeneous prevalence of perinatal depression, with the review's authors explicitly flagging social determinants — isolation, reduced access to services, and constrained decision-making autonomy — as central rather than incidental to the elevated rural burden [6].




Figure 1: Odds ratios for the three factors most strongly associated with postpartum depression in a pooled meta-analysis of 11 Ethiopian studies (n=7,582) -- poor social support carries the largest effect, and is also the risk factor most directly targeted by psychosocial empowerment programs [4].

What is striking about the risk-factor literature, and what motivates the empowerment framing specifically, is how consistently the strongest predictors of postpartum depression are psychosocial rather than purely biological or obstetric. Zeleke and colleagues' finding that poor social support carries a larger effect size than either domestic violence or unplanned pregnancy is a useful anchor point: of the three major risk factors identified, social support is also the one most directly and plausibly modifiable by intervention, since a clinician cannot retroactively make a pregnancy planned, and addressing domestic violence requires an

entirely different, protection-oriented intervention architecture, but building social support infrastructure around a new mother is squarely within the scope of what a psychosocial program can attempt to do [4]. O'Hara and McCabe's authoritative review of the postpartum depression literature similarly emphasizes that while biological vulnerability (prior depression history, hormonal sensitivity) sets a baseline risk, the proximal triggers that determine whether that vulnerability is expressed clinically are overwhelmingly psychosocial: the quality of the marital or partner relationship, the adequacy of practical and emotional support, and the degree of stress associated with infant care and role transition [7].

This is precisely the gap that empowerment-oriented psychosocial interventions are designed to close, and it is worth being specific about the causal chain such interventions implicitly claim: that increasing a woman's sense of control and competence over her postpartum experience, connecting her to peer or community support, and providing structured coping skills will reduce the psychosocial risk exposure (isolation, felt helplessness, unaddressed stress) that the epidemiological literature identifies as the dominant modifiable driver of postpartum depressive symptoms. Garcia and Yim's systematic review of empowerment-related constructs in the perinatal period found that measures of empowerment — spanning autonomy, self-efficacy, and social support — were consistently, if heterogeneously, associated with lower perinatal depressive symptom burden and, in some studies, with reduced risk of premature birth, across a literature that the authors note suffers from considerable conceptual fragmentation (different studies operationalize "empowerment" quite differently) but converges on a broadly consistent directional finding [8].

3. Empowerment as a Theoretical Framework

Before assessing whether empowerment-based programs work, it is worth being precise about what "empowerment" means as an intervention target, because the term is used loosely enough in both clinical and lay contexts that it risks becoming an empty signifier. Zimmerman's foundational framework, developed originally in community psychology rather than maternal health specifically, distinguishes three interlocking components: intrapersonal empowerment, the psychological sense that one has control and

competence over one's circumstances; interactional empowerment, the practical understanding and skills needed to actually navigate one's social and institutional environment (knowing how to access a lactation consultant, how to advocate with a pediatrician, how to set a boundary with an unhelpful relative); and behavioral empowerment, the concrete actions a person takes as a result of the first two [2]. This tripartite structure matters for evaluating postpartum interventions because a program can plausibly succeed at building intrapersonal empowerment (a mother feels more confident) without translating that into interactional or behavioral empowerment (she still does not know how or feel able to access support when she actually needs it), and the postpartum-specific literature suggests that programs targeting all three components tend to outperform those targeting confidence alone.

Bandura's social cognitive theory offers a complementary and more mechanistically specific account, centered on self-efficacy: the belief that one is capable of executing a specific behavior needed to produce a desired outcome, as distinct from a generalized sense of confidence [3]. Bandura's framework identifies four sources of self-efficacy — mastery experiences (successfully doing the thing), vicarious experience (watching someone similar to you do it successfully), verbal persuasion (being told, credibly, that you can do it), and physiological/affective states (interpreting your own anxiety as manageable rather than disqualifying) — and this taxonomy maps unusually cleanly onto the mechanics of postpartum psychosocial programs. Peer support interventions work primarily through vicarious experience and verbal persuasion (a peer who has been through postpartum depression herself is a uniquely credible source of both); structured self-management and skills-training programs work primarily through mastery experience (practicing a coping technique and seeing it work); and group-based care models work through some combination of all four, since they provide social modeling, direct skill practice, and a validating, symptom-normalizing social context simultaneously.

The empirical postpartum-specific literature has, for the most part, adopted this theoretical machinery directly rather than developing bespoke maternal-mental-health theory from scratch. A 2026 randomized trial of a self-management empowerment program for

cesarean-delivery mothers, for instance, explicitly framed its intervention around increasing sense of coherence (a related construct describing the degree to which one experiences the world as comprehensible, manageable, and meaningful) and self-efficacy as the mechanistic pathway to reduced postpartum anxiety, and found significant improvements in both the mediating constructs and the anxiety outcome itself among women who received the structured empowerment program relative to standard care [9]. This kind of explicit mechanism-testing, where a trial measures not just the downstream mental health outcome but also the theorized psychological mediator, is still relatively uncommon in this literature but represents, in our view, the methodological direction the field most needs more of, because it is the only way to distinguish a program that works from a program that works for the reason its designers think it does.

4. Peer Support and Community-Delivered Empowerment Programs

The most extensively trialed and most consistently positive strand of psychosocial empowerment intervention for postpartum mental health is peer support: structured programs in which women who have themselves experienced postpartum depression, or who are simply trained community members, provide support — emotional, informational, and practical — to new mothers, typically through a combination of telephone contact, home visits, or group sessions. This model is attractive on empowerment-theoretic grounds precisely because it operationalizes Bandura's vicarious-experience and verbal-persuasion pathways directly: the peer is a credible, relatable model of recovery, and her encouragement carries weight that a clinician's more distant reassurance may not [3].

The single most influential body of work in this space comes from Atif Rahman and colleagues' Thinking Healthy Programme, originally developed and tested in rural Pakistan. The foundational 2008 cluster-randomized controlled trial delivered a cognitive-behaviorally grounded intervention through trained community health workers — not specialist mental health professionals — to depressed mothers in their own homes, and found substantial reductions in depression at both the immediate post-intervention assessment and at 12-month follow-up relative to enhanced usual care, alongside improvements in

maternal functioning and several child health outcomes [10]. The trial's significance extends well beyond its effect size: it demonstrated that a structured psychosocial intervention could be delivered by non-specialist community workers at meaningful scale in a low-resource rural setting, which reframed maternal mental health intervention from a specialist-clinic problem into a task-shareable, community-deliverable one. A 2015 long-term follow-up of the same cohort found that the intervention's benefits extended to child cognitive and socioemotional development outcomes measured years after the original trial, evidence that maternal psychosocial recovery has developmental consequences for children that persist well beyond the postpartum window itself [11].

Table 1: Representative trials and meta-analyses of peer-support and community-delivered psychosocial empowerment interventions for postpartum mental health, spanning delivery model, population, and reported effect.

Study	Delivery Model / Population	Design	Reported Effect
Rahman et al. [10]	CHW-delivered CBT-based home visits, rural Pakistan	Cluster-RCT	Substantial depression reduction at post-intervention and 12-month follow-up vs. enhanced usual care
Sikander et al. [12]	Thinking Healthy Programme, trained volunteer peers, Pakistan	Cluster-RCT	Meaningful effectiveness retained with peer (non-specialist) delivery model
Boran et al. [13]	Thinking Healthy Programme, universal group format in routine antenatal care, Turkey	Randomized pilot	Feasible and acceptable integration into routine antenatal care
Nisar et al. [14]	WHO Thinking Healthy Programme, routine antenatal care, China	Randomized pilot	Feasible cross-cultural integration outside original South Asian setting
Huang et al. [15]	Peer support, perinatal depression (pooled trials)	Systematic review & meta-analysis	Significant pooled reduction in perinatal depressive symptoms vs. control
Fang et al. [16]	Peer support, perinatal depression (pooled trials)	Meta-analysis	Significant beneficial effect on perinatal depression, independently convergent with Huang et al.
Shorey et al. [17]	Technology (app)-mediated peer support for postnatal depression prevention	RCT	Meaningful benefit vs. standard care with app-mediated delivery

The Thinking Healthy Programme has since been adapted and re-tested across multiple health systems and delivery models, and this adaptation history is itself informative about which components of the intervention are portable and which are context-dependent. A 2019 cluster-randomized trial tested delivery specifically through trained volunteer peers rather than paid community health workers, finding that the peer-delivered model retained meaningful effectiveness, which matters directly for sustainability and cost in low-resource settings where paid specialist staffing is the binding constraint [12]. More recent pilot trials have tested the programme as a universal, group-based intervention integrated into routine antenatal care in Turkey, finding it feasible and acceptable within an existing health system rather than as a stand-alone specialist referral pathway [13], and a parallel pilot integrated the WHO-endorsed programme into routine antenatal care in China, extending the evidence base for this model beyond its South Asian origins into an entirely different health system and cultural context [14]. Taken together, this sequence of trials — from a single-country efficacy trial through delivery-model variation to cross-cultural adaptation and integration into routine care — is close to the ideal evidentiary trajectory an intervention needs to travel before it can be recommended as a general-purpose model rather than a context-specific finding, and the Thinking Healthy Programme is, in our reading, further along that trajectory than any other psychosocial empowerment intervention for postpartum mental health currently in the literature.

Two recent meta-analyses provide the clearest population-level synthesis of the peer support literature as a whole, independent of any single program brand. Huang and colleagues' 2020 systematic review and meta-analysis found that peer support interventions produced a statistically significant reduction in perinatal depressive symptoms relative to control conditions across the pooled trial evidence, alongside favorable findings on cost and participant satisfaction, though the authors also noted considerable heterogeneity in intervention format (telephone-based, in-person, group versus individual) across the included trials [15]. Fang and colleagues' 2022 meta-analysis reached a broadly convergent conclusion, finding a significant beneficial effect of peer support on perinatal depression outcomes across the studies synthesized, and, notably, both

meta-analyses were conducted independently using different inclusion criteria and largely non-overlapping study sets, which strengthens confidence that the peer-support effect is not an artifact of any single research group's methodology or publication practices [16]. Technology has increasingly mediated how this peer support is delivered: a 2019 randomized controlled trial by Shorey and colleagues evaluated a technology-based (mobile app-mediated) peer support program aimed at preventing postnatal depression and found meaningful benefit for participants receiving the intervention relative to standard care, evidence that the underlying peer-support mechanism generalizes reasonably well from in-person and telephone delivery to app-mediated delivery, which matters considerably for scalability [17].

It is worth pausing on why social support specifically — the mechanism peer support interventions are most directly targeting — carries such epidemiological weight. Khademi and Kaveh's 2024 systematic review and logic-framework synthesis of social support as a coping resource in the postpartum period found that social support operates through multiple simultaneous pathways: it buffers the direct stress of infant care, it provides practical assistance that reduces the objective burden of caregiving tasks, and it validates and normalizes the emotional difficulty of the postpartum transition in a way that reduces a mother's sense of isolated deficiency [18]. This multi-pathway account helps explain why peer support interventions, despite being comparatively low-cost and low-intensity relative to individual psychotherapy, have produced effect sizes in the meta-analytic literature that are clinically meaningful rather than marginal: they are not a diluted substitute for more intensive intervention but a mechanistically well-matched one, targeting precisely the risk factor — poor social support — that carries the largest effect size in factor-analytic studies of postpartum depression risk [4].

5. Group Prenatal Care and Structural Empowerment Models

A structurally distinct approach to psychosocial empowerment reorganizes the delivery of routine prenatal and postnatal care itself around group process, rather than adding a separate empowerment-focused program on top of standard individual visits. The best-known model of this kind is CenteringPregnancy,

which replaces individual prenatal visits with facilitated group sessions in which women at similar gestational stages receive clinical assessment alongside structured facilitated discussion, peer interaction, and skill-building, explicitly designed around a relationship-centered rather than purely biomedical care philosophy [19]. The empowerment logic of this model differs somewhat from the peer-support literature discussed above: rather than adding a peer-support layer to standard individual care, it restructures the care encounter itself so that social connection, mutual learning, and active participation are built into every visit rather than delivered as a supplementary intervention.

The trial evidence on group prenatal care's psychosocial effects is smaller than the peer-support literature but has grown substantially in the past decade. Heberlein and colleagues' 2016 study comparing group and individual prenatal care found measurable differences in psychosocial outcomes favoring the group model, and a subsequent 2023 randomized controlled trial by Smith and colleagues directly compared group versus individual prenatal care on psychosocial outcomes with a more rigorous experimental design, adding needed causal evidence to a literature that had previously relied more heavily on observational comparison [20,21]. A 2023 UK pilot randomized controlled trial of a group antenatal care model (REACH Pregnancy Circles) similarly found the group format feasible and acceptable, extending this evidence base outside the U.S. health system in which CenteringPregnancy was originally developed and most extensively tested [22]. Most recently, a 2026 randomized controlled trial by Melnyk and colleagues tested a group prenatal care intervention specifically targeting postpartum mental health and healthy lifestyle behaviors among women of color experiencing depression, anxiety, and stress during pregnancy — a population that carries disproportionate psychosocial risk burden and has been comparatively underrepresented in the earlier group-care trial literature — and found improvements in postpartum mental health outcomes among participants [23]. The explicit focus on a population facing compounded structural and psychosocial risk marks a meaningful maturation of this literature beyond its earlier, largely demographically homogeneous trial base.

There is a reasonable theoretical case, though the direct

comparative trial evidence remains thinner than one would like, that group-based structural models and individually-delivered peer support models are not competing approaches but complementary ones addressing different components of Zimmerman's tripartite empowerment framework [2]. Group prenatal care builds interactional empowerment directly into the care relationship itself — women practice asking questions, comparing experiences, and building a shared vocabulary for their concerns within the clinical encounter — while dedicated peer support programs, particularly those extending into the postpartum period specifically rather than concentrating on the antenatal window, more directly target the intrapersonal and behavioral empowerment components through modeling and encouragement from someone who has already navigated the specific challenge of postpartum recovery [10,12]. Programs that combine structural group care during pregnancy with continued peer support after birth, rather than treating the perinatal period as ending at delivery, would on this theoretical account be expected to outperform either component alone, though we are not aware of a trial specifically designed to test this combined-model hypothesis head-to-head against either component in isolation.

6. Empowerment Programs for Compounded Psychosocial Risk

Empowerment language carries particular clinical and ethical weight when applied to women facing intimate partner violence during pregnancy or the postpartum period, both because domestic violence is itself among the strongest predictors of postpartum depression identified in the epidemiological literature [4] and because interventions in this space must be designed with safety, not just symptom reduction, as a primary outcome. Martinez-Torteya and colleagues' 2025 pilot randomized controlled trial of the Pregnant Moms' Empowerment Program, adapted and tested in Mexico specifically for women exposed to intimate partner violence, found that the program produced measurable improvements in mental health outcomes for participants, evidence that a structured empowerment-based intervention can be both safe and effective for this higher-risk, higher-complexity population rather than requiring an entirely separate intervention architecture [24]. This finding is clinically significant because it addresses a legitimate concern that has historically limited psychosocial intervention

research with IPV-exposed populations — namely, that standard supportive or peer-based interventions might be inappropriate or even unsafe for women navigating an ongoing abusive relationship — by demonstrating feasibility and benefit within a trial explicitly designed with this population's safety needs in mind.

The relationship between compounded psychosocial risk and mother-infant relational outcomes deserves separate attention, because postpartum depression's consequences extend beyond the mother's own symptom burden into the earliest mother-infant relationship, and empowerment-oriented interventions are increasingly being evaluated on this dyadic outcome as well as on maternal symptoms alone. Badr and colleagues' 2018 study examining whether the effect of postpartum depression on mother-infant bonding is consistent across different sociocultural contexts found that while the depression-bonding association is broadly observed, its magnitude and the specific mechanisms through which it operates vary meaningfully by context, which argues against treating any single psychosocial intervention model as a universal solution without local adaptation [25]. Kornfield and colleagues' 2021 study of risk and resilience factors influencing postpartum depression and mother-infant bonding during the COVID-19 pandemic offers a particularly instructive natural experiment: the pandemic sharply reduced the availability of precisely the in-person social support and peer contact that the interventions discussed throughout this article are designed to provide, and the study found that reduced access to social support during this period was associated with worse maternal mental health and bonding outcomes, which is indirect but reasonably strong evidence for the causal importance of the social-support mechanism these programs target, observed under conditions where the intervention (ordinary social support) was largely involuntarily withdrawn rather than experimentally withheld [26].

7. Mechanisms, Effect Sizes, and What the Evidence Actually Shows

It is worth stepping back from the individual-trial narrative to ask plainly what the accumulated evidence supports and what it does not. On the positive side, the case for psychosocial empowerment programs reducing postpartum depressive and anxiety symptoms is considerably stronger than the case for most complementary or alternative postpartum mental health

interventions: it rests on two independent, non-overlapping meta-analyses of the peer-support literature specifically [15,16], a well-replicated and cross-culturally adapted flagship program (the Thinking Healthy Programme) with trial evidence spanning Pakistan, Turkey, and China and multiple delivery models [10,12,13,14], a growing and increasingly methodologically rigorous group prenatal care literature [20,21,23], and at least one well-designed trial extending the evidence base to a compounded-risk population that earlier research had largely excluded [24]. The Cochrane systematic reviews covering this space, though somewhat dated in their most recent full updates, similarly conclude that psychosocial and psychological interventions carry a genuine, if moderate, protective and therapeutic effect on postpartum depression, a conclusion broadly consistent with the more recent individual trials and meta-analyses synthesized here [27,28].

On the more cautious side, several patterns recur across this literature that temper how confidently it should be generalized. First, effect sizes across the peer-support meta-analyses, while statistically significant and clinically non-trivial, are moderate rather than large, and neither major meta-analysis found peer support sufficient to fully resolve depression for the majority of moderately-to-severely symptomatic participants — these are best understood as effective preventive and adjunctive interventions rather than replacements for clinical treatment in more severe cases [15,16]. Second, intervention fidelity and format vary enormously across the included studies — telephone-based, in-person, app-mediated, individual, and group delivery are all represented under the single label "peer support," and the field has not yet cleanly established which specific format features (session frequency, peer training intensity, duration of contact) drive the observed effect, which limits how precisely a new implementation can be designed from the existing evidence base. Third, and this echoes a critique familiar from other psychosocial intervention literatures, the great majority of trials measure symptom reduction on standardized screening instruments like the EPDS [1] without directly measuring the theorized empowerment mechanism itself (perceived control, self-efficacy, social support quality) as a mediator, which means that while the theoretical case for why these programs should work is strong, direct mechanistic confirmation — showing that the

intervention changed the mediator and the mediator change explains the outcome change — remains the exception rather than the rule in this literature, with the self-management empowerment trial among cesarean mothers standing out as a rarer example of an explicitly mediation-tested design [9].

Fourth, generalizability across settings remains uncertain despite the encouraging cross-cultural adaptation history of the Thinking Healthy Programme. Badr and colleagues' finding that the depression-bonding relationship varies by sociocultural context is a useful caution against assuming an intervention validated in one setting transfers with equal effectiveness elsewhere without local adaptation and re-testing [25]. Finally, most trials in this literature are comparatively short in follow-up duration; the Thinking Healthy Programme's 2015 developmental follow-up study is a valuable exception precisely because it is uncommon, and the field would benefit from more trials tracking outcomes well beyond the immediate postpartum window [11].

8. Open Challenges

Several open problems recur across this literature and are worth naming directly rather than leaving implicit. First, as noted above, mechanistic testing remains underdeveloped: the field has strong theoretical grounds, from both Zimmerman's empowerment framework and Bandura's self-efficacy theory, for expecting these interventions to work through increased perceived control and social connectedness, but relatively few trials directly measure and test these mediating pathways rather than simply reporting the downstream symptom outcome [2,3,9]. Second, dose-response and active-ingredient questions remain largely unanswered: no trial we are aware of has systematically varied peer-contact frequency, session duration, or facilitator training intensity within a single study design to identify which components of a multi-component empowerment program are load-bearing and which are supplementary, which matters considerably for cost-effective scale-up in resource-constrained settings.

Third, the compounded-risk literature, while growing, remains thin relative to its clinical importance: intimate partner violence is among the strongest psychosocial risk factors identified in the epidemiological literature [4], yet trial evidence specifically evaluating

empowerment interventions for IPV-exposed postpartum populations is limited essentially to a single well-designed pilot study at the time of this synthesis [24], and a pilot trial, however well-conducted, cannot substitute for a larger, adequately powered confirmatory trial before this model can be recommended at scale. Fourth, most of the group prenatal care literature and much of the peer-support literature have historically underrepresented racial and ethnic minority populations and populations facing intersecting structural disadvantage, a gap the 2026 Melnyk trial specifically targeting women of color begins to address but does not close on its own [23]. Fifth, digital and app-mediated delivery models, while showing encouraging early results in the Shorey trial [17], have not yet been tested at the scale or with the follow-up duration needed to know whether technology-mediated peer support produces durable benefit comparable to in-person delivery, or whether it functions best as a genuine substitute versus a supplementary access channel for women who cannot attend in-person programming.

Sixth, outcome measurement itself is less standardized across this literature than it first appears. Most trials rely on the Edinburgh Postnatal Depression Scale as the primary outcome instrument [1], which is a real methodological strength in that it allows some degree of cross-study comparison, but the threshold used to define a "case" of clinically significant depression varies from study to study, as does the timing of administration relative to delivery, and a meta-analysis that pools studies using different cutoffs and different assessment windows is, to some degree, pooling somewhat different constructs under a single label. Anxiety outcomes are measured even less consistently, with some trials using dedicated anxiety instruments and others relying on the EPDS's anxiety-adjacent items as a proxy, which likely understates the true effect of empowerment interventions on anxiety specifically, since anxiety and depressive symptoms in the postpartum period are highly comorbid but not identical constructs [5]. A more consistent core outcome set across this literature, analogous to core outcome sets that have improved comparability in other areas of clinical trial research, would make future meta-analytic synthesis considerably more informative than it currently is.

Finally, and more structurally, the field has not

adequately reconciled the individual-level empowerment framing common in this literature with the fact that several of the strongest risk factors for postpartum depression — poor social support, domestic violence, unplanned pregnancy — are themselves substantially shaped by structural conditions (housing stability, economic security, availability of accessible healthcare, the presence or absence of legal and social protections against intimate partner violence) that an individually-targeted psychosocial program, however well-designed, cannot fully address on its own [4]. This is not an argument against psychosocial empowerment programs, which the evidence reviewed here supports as genuinely effective within their scope; it is an argument for locating them honestly within a broader public health response rather than treating individual-level empowerment intervention as a sufficient answer to what is, in significant part, a structurally produced problem.

9. Conclusion

The accumulated evidence, read as a whole, supports a fairly specific and moderately confident claim: psychosocial empowerment programs — peer support, community health worker-delivered structured psychotherapy, group-based prenatal and postnatal care, and targeted self-management programs — produce a real, meaningful, and now multiply-replicated reduction in postpartum depressive and anxiety symptoms, grounded in a coherent theoretical account of why they should work (Zimmerman's empowerment framework and Bandura's self-efficacy theory) and increasingly corroborated by direct trial evidence measuring not just symptom outcomes but, in a growing number of studies, the psychosocial mediators these theories predict [2,3,9]. The Thinking Healthy Programme's trajectory from a single rural Pakistani trial to a cross-culturally adapted, multiply-replicated model integrated into routine antenatal care across several health systems is, in our reading, the clearest existence proof that this class of intervention is not a narrow, context-bound curiosity but a genuinely portable model for task-shared, community-deliverable maternal mental health care [10,12,13,14].

At the same time, the evidence base has real and specific limits that future work should target directly: sparse mechanistic testing of the empowerment pathway itself, limited dose-response and component-isolation

research, thin trial coverage of intimate-partner-violence-exposed and other compounded-risk populations relative to their epidemiological importance, uncertain durability and generalizability of digital delivery models, and an underdeveloped connection between individual-level empowerment intervention and the structural conditions that produce much of the psychosocial risk these programs are trying to counteract. None of these gaps undermine the core finding that psychosocial empowerment approaches work; they define, instead, the agenda for making this already promising class of intervention more precisely targeted, more equitably distributed, and more honestly positioned within the larger set of structural and clinical responses that postpartum mental health, as a genuinely biopsychosocial condition, actually requires.

10. References

1. Cox, J.L., Holden, J.M., Sagovsky, R. (1987) Detection of Postnatal Depression: Development of the 10-Item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry*, 150(6): 782-786. DOI: [10.1192/bjp.150.6.782](https://doi.org/10.1192/bjp.150.6.782)
2. Zimmerman, M.A. (1995) Psychological Empowerment: Issues and Illustrations. *American Journal of Community Psychology*, 23(5): 581-599. DOI: [10.1007/bf02506983](https://doi.org/10.1007/bf02506983)
3. Bandura, A. (2004) Health Promotion by Social Cognitive Means. *Health Education & Behavior*, 31(2): 143-164. DOI: [10.1177/1090198104263660](https://doi.org/10.1177/1090198104263660)
4. Zeleke, T.A., Getinet, W., Tessema, Z.T., Gebeyehu, K. (2021) Prevalence and Associated Factors of Post-Partum Depression in Ethiopia: A Systematic Review and Meta-Analysis. *PLOS ONE*, 16(2): e0247005. DOI: [10.1371/journal.pone.0247005](https://doi.org/10.1371/journal.pone.0247005)
5. Ou, L., Shen, Q., Xiao, M., Wang, W., He, T., Wang, B. (2025) Prevalence of Co-Morbid Anxiety and Depression in Pregnancy and Postpartum: A Systematic Review and Meta-Analysis. *Psychological Medicine*, 55: e84. DOI: [10.1017/s0033291725000601](https://doi.org/10.1017/s0033291725000601)
6. Pan, T., Zeng, Y., Chai, X., Wen, Z., Tan, X., Sun, M. (2024) Global Prevalence of Perinatal Depression and Its Determinants Among Rural Women: A Systematic Review and Meta-Analysis. *Depression and Anxiety*, 2024: 1882604. DOI: [10.1155/2024/1882604](https://doi.org/10.1155/2024/1882604)
7. O'Hara, M.W., McCabe, J.E. (2013) Postpartum Depression: Current Status and Future Directions. *Annual Review of Clinical Psychology*, 9(1): 379-407. DOI: [10.1146/annurev-clinpsy-050212-185612](https://doi.org/10.1146/annurev-clinpsy-050212-185612)
8. Garcia, E.R., Yim, I.S. (2017) A Systematic Review of Concepts Related to Women's Empowerment in the Perinatal Period and their Associations with Perinatal Depressive Symptoms and Premature Birth. *BMC Pregnancy and Childbirth*, 17(S2): 347. DOI: [10.1186/s12884-017-1495-1](https://doi.org/10.1186/s12884-017-1495-1)
9. Najafi ShahaliBegloo, F., Kamali, F., Mohamadi, S., Ravanipour, M. (2026) Assessment of Self-Management Empowerment Program on Sense of Coherence, Self-Efficacy and Postpartum Anxiety in Cesarean Mothers: Randomized Trial. *Scientific Reports*, 16(1): 7975. DOI: [10.1038/s41598-026-37944-7](https://doi.org/10.1038/s41598-026-37944-7)
10. Rahman, A., Malik, A., Sikander, S., Roberts, C., Creed, F. (2008) Cognitive Behaviour Therapy-Based Intervention by Community Health Workers for Mothers with Depression and their Infants in Rural Pakistan: A Cluster-Randomised Controlled Trial. *The Lancet*, 372(9642): 902-909. DOI: [10.1016/s0140-6736\(08\)61400-2](https://doi.org/10.1016/s0140-6736(08)61400-2)

11. Maselko, J., Sikander, S., Bhalotra, S., Bangash, O., Ganga, N., Mukherjee, S., et al. (2015) Effect of an Early Perinatal Depression Intervention on Long-Term Child Development Outcomes: Follow-Up of the Thinking Healthy Programme Randomised Controlled Trial. *The Lancet Psychiatry*, 2(7): 609-617. DOI: [10.1016/s2215-0366\(15\)00109-1](https://doi.org/10.1016/s2215-0366(15)00109-1)
12. Sikander, S., Ahmad, I., Atif, N., Zaidi, A., Vanobberghen, F., Weiss, H.A., et al. (2019) Delivering the Thinking Healthy Programme for Perinatal Depression Through Volunteer Peers: A Cluster Randomised Controlled Trial in Pakistan. *The Lancet Psychiatry*, 6(2): 128-139. DOI: [10.1016/s2215-0366\(18\)30467-x](https://doi.org/10.1016/s2215-0366(18)30467-x)
13. Boran, P., Dönmez, M., Barış, E., Us, M.C., Altaş, Z.M., Nisar, A., et al. (2023) Delivering the Thinking Healthy Programme as a Universal Group Intervention Integrated into Routine Antenatal Care: A Randomized-Controlled Pilot Study. *BMC Psychiatry*, 23(1): 14. DOI: [10.1186/s12888-022-04499-6](https://doi.org/10.1186/s12888-022-04499-6)
14. Nisar, A., Yin, J., Zhang, J., Qi, W., Yu, J., Li, J., Li, X., Rahman, A. (2025) Integrating WHO Thinking Healthy Programme for Maternal Mental Health into Routine Antenatal Care in China: A Randomized-Controlled Pilot Trial. *Frontiers in Global Women's Health*, 5: 1475430. DOI: [10.3389/fgwh.2024.1475430](https://doi.org/10.3389/fgwh.2024.1475430)
15. Huang, R., Yan, C., Tian, Y., Lei, B., Yang, D., Liu, D., Lei, J. (2020) Effectiveness of Peer Support Intervention on Perinatal Depression: A Systematic Review and Meta-Analysis. *Journal of Affective Disorders*, 276: 788-796. DOI: [10.1016/j.jad.2020.06.048](https://doi.org/10.1016/j.jad.2020.06.048)
16. Fang, Q., Lin, L., Chen, Q., Yuan, Y., Wang, S., Zhang, Y., Liu, T., Cheng, H., Tian, L. (2022) Effect of Peer Support Intervention on Perinatal Depression: A Meta-Analysis. *General Hospital Psychiatry*, 74: 78-87. DOI: [10.1016/j.genhosppsych.2021.12.001](https://doi.org/10.1016/j.genhosppsych.2021.12.001)
17. Shorey, S., Chee, C.Y.I., Ng, E.D., Lau, Y., Dennis, C.L., Chan, Y.H. (2019) Evaluation of a Technology-Based Peer-Support Intervention Program for Preventing Postnatal Depression (Part 1): Randomized Controlled Trial. *Journal of Medical Internet Research*, 21(8): e12410. DOI: [10.2196/12410](https://doi.org/10.2196/12410)
18. Khademi, K., Kaveh, M.H. (2024) Social Support as a Coping Resource for Psychosocial Conditions in Postpartum Period: A Systematic Review and Logic Framework. *BMC Psychology*, 12(1): 301. DOI: [10.1186/s40359-024-01814-6](https://doi.org/10.1186/s40359-024-01814-6)
19. Massey, Z., Rising, S.S., Ickovics, J. (2006) CenteringPregnancy Group Prenatal Care: Promoting Relationship-Centered Care. *Journal of Obstetric, Gynecologic & Neonatal Nursing*, 35(2): 286-294. DOI: [10.1111/j.1552-6909.2006.00040.x](https://doi.org/10.1111/j.1552-6909.2006.00040.x)
20. Heberlein, E.C., Picklesimer, A.H., Billings, D.L., Covington-Kolb, S., Farber, N., Frongillo, E.A. (2016) The Comparative Effects of Group Prenatal Care on Psychosocial Outcomes. *Archives of Women's Mental Health*, 19(2): 259-269. DOI: [10.1007/s00737-015-0564-6](https://doi.org/10.1007/s00737-015-0564-6)
21. Smith, J.C., Heberlein, E.C., Domingue, A., LaBoy, A., Britt, J., Crockett, A.H. (2023) Randomized Controlled Trial on the Effect of Group Versus Individual Prenatal Care on Psychosocial Outcomes. *Journal of Obstetric, Gynecologic & Neonatal Nursing*, 52(6): 467-480. DOI: [10.1016/j.jogn.2023.07.006](https://doi.org/10.1016/j.jogn.2023.07.006)
22. Sawtell, M., Wiggins, M., Wiseman, O., et al. (2023) Group Antenatal Care: Findings from a Pilot Randomised Controlled Trial of REACH Pregnancy Circles. *Pilot and Feasibility Studies*, 9(1): 91. DOI: [10.1186/s40814-023-01238-w](https://doi.org/10.1186/s40814-023-01238-w)
23. Melnyk, B.M., Gennaro, S., Szalacha, L.A., Gibeau, A.M., Hoying, J., O'Connor, C.M., Cooper, A., Aviles, M.M. (2026) Improving Postpartum Mental Health and Healthy Lifestyle Behaviors in Women of Color Who Are Pregnant and Experiencing Depression, Anxiety, and Stress Through Group Prenatal Care: A Randomized Controlled Trial. *Journal of Midwifery & Women's Health*. DOI: [10.1111/jmwh.70154](https://doi.org/10.1111/jmwh.70154)
24. Martinez-Torteya, C., Miller-Graff, L.E., Carney, J.R., Esparza-Dávila, S.P., Acuapa-Juárez, J.C., Howell, K.H. (2025) The Pregnant Moms' Empowerment Program - Mexico Enhances Mental Health for Women Exposed to Intimate Partner Violence: A Pilot Randomized Controlled Trial. *Archives of Women's Mental Health*, 28(4): 947-959. DOI: [10.1007/s00737-024-01549-8](https://doi.org/10.1007/s00737-024-01549-8)
25. Badr, L.K., Ayyazian, N., Lameh, S., Charafeddine, L. (2018) Is the Effect of Postpartum Depression on Mother-Infant Bonding Universal? *Infant Behavior and Development*, 51: 15-23. DOI: [10.1016/j.infbeh.2018.02.003](https://doi.org/10.1016/j.infbeh.2018.02.003)
26. Kornfield, S.L., White, L.K., Waller, R., Njoroge, W., Barzilay, R., et al. (2021) Risk and Resilience Factors Influencing Postpartum Depression and Mother-Infant Bonding During COVID-19. *Health Affairs*, 40(10): 1566-1574. DOI: [10.1377/hlthaff.2021.00803](https://doi.org/10.1377/hlthaff.2021.00803)
27. Dennis, C.L., Dowswell, T. (2013) Psychosocial and Psychological Interventions for Preventing Postpartum Depression. *Cochrane Database of Systematic Reviews*, 2013(2): CD001134. DOI: [10.1002/14651858.cd001134.pub3](https://doi.org/10.1002/14651858.cd001134.pub3)
28. Dennis, C.L., Hodnett, E. (2006) Psychosocial and Psychological Interventions for Treating Postpartum Depression. *Cochrane Database of Systematic Reviews*, 2006(4): CD006116. DOI: [10.1002/14651858.cd006116](https://doi.org/10.1002/14651858.cd006116)

11. Declarations

Funding

This literature synthesis received no dedicated grant funding from public, commercial, or not-for-profit funding bodies.

Conflict of Interest

The authors report no competing interests related to this work.

Data Availability Statement

This article synthesizes previously published literature and generated no new primary data; readers are directed to the cited sources for their respective data and materials.

Ethical Approval

Not applicable, as this review involved no human participants, animal subjects, or original data collection.

Citation: Fernandez M, Salgado P, Valdes R. "Psychosocial Empowerment Programs and Their Effect on Postpartum Mental Health." J Women Health Saf Empower (2026): 001-012 doi: [10.5281/zenodo.22849484](https://doi.org/10.5281/zenodo.22849484)